



# Fact Sheet

## Postemergence broadleaf weed control in field corn

### Product Advantages:

- Can be applied up to 20 inch tall corn
- Two modes of action for resistance management
- Minimal concerns for volatilization or drift
- Excellent crop safety
- Wide application window
- Does not contain atrazine or 2,4-D

### Weed spectrum:

Canada thistle <sup>1</sup>	Lambsquarters <sup>1</sup>	Russian thistle <sup>1</sup>	Wild buckwheat <sup>1</sup>
Cocklebur	Nightshades <sup>1</sup>	Shepard's purse	Wild mustard
Horseweed	Pigweed <sup>1</sup>	Smartweed	Velvetleaf
Jimsonweed	Purslane	Sunflower	Wormwood, biennial
Kochia <sup>1</sup>	Ragweeds	Velvetleaf	<sup>1</sup> partial control (weeds less than 2" tall)

### Application timing:

- Postemergence - Broadcast up to 20 inch tall corn
- Target broadleaves should be less than 6 inches tall (see label for rates)

### Herbicide Components:

- 1.5 ounces (a.e.) clopyralid + .55 ounce (a.i.) flumetsulam (3.0 oz/acre rate)

### Mixing instructions:

- 2 - 5 ounces per acre (consult label for recommended rates)
- Add NIS at 1 quart per 100 gallons or COC at 1 gallon per 100 gallons
- Use 10 to 40 gallons of water, adjust pressure to control drift
- Under dry conditions, add UAN at 2.5 gallons per 100 gallons to enhance uptake
- Do not use undiluted UAN as total carrier
- Rainfast in 6 hours

### Other Information:

- PPE: goggles, waterproof gloves, long pants, long sleeved shirt, shoes and socks
  - Not a Restricted Use Product, REI is 48 hours
  - Do not apply Hornet if corn was previously treated with Counter or Thimet
  - Crop Rotational Interval:
- |                              |                       |
|------------------------------|-----------------------|
| oats, wheat, barley, rye     | 4 months              |
| alfalfa, soybeans, dry beans | 10.5 months           |
| grain sorghum                | 12 months             |
| potatoes, peas, sunflowers   | 18 months             |
| sugar beets, canola          | 26 months & bio-assay |