



Rapport™ BroadSpec Herbicide

Rapport™ BroadSpec Selective Herbicide contains thifensulfuron and tribenuron, the active ingredients used in Affinity™ BroadSpec.



Rapport™ BroadSpec for Fast Control of Tough Weeds

Tough weeds like **Canada thistle** and **wild buckwheat** can take over your wheat crop and **cut into your bottom line**. Rapport™ BroadSpec provides excellent control of Canada thistle, wild buckwheat and a broad spectrum of other difficult weeds. Plus Rapport BroadSpec gives you rotational flexibility for next season.

KEY PERFORMANCE BENEFITS

Dependable post-emergent control of a broad spectrum of broadleaf weeds
Improved Canada thistle control
Safe to the growing crop
Excellent tank-mix partner
Flexible rotation options

ACTIVE INGREDIENTS

thifensulfuron-methyl
and tribenuron-methyl

% ACTIVE INGREDIENTS

25% and 25%

FORMULATION

dry flowable

CHEMICAL FAMILY

sulfonylurea

GROUP NUMBER

2

EPA REGISTRATION

71368-79

BRAND ALTERNATIVE

Affinity™ BroadSpec

MODE OF ACTION

ALS inhibitor

RESTRICTED USE

no

SIGNAL WORD

caution

PACKAGING INFORMATION

8 x 48 oz



LABELED USES

Wheat
Barley
Fallow

WEEDS CONTROLLED

Annual knawel	Marestail**
Annual sowthistle	Marshelder
Black mustard	Mayweed chamomile
Blue/purple mustard	Miners lettuce
Broadleaf dock	Narrowleaf lambquarters
Bur buttercup	Nightflowering catchfly
Bushy wallflower/ treacle mustard	Nightshade (<i>cutleaf</i> , <i>hairy</i>)**
Canada thistle*	Pennsylvania smartweed
Catchweed bedstraw**	Pineappleweed
Clasping pepperweed	Prickly lettuce*
Coast fiddleneck	Prostrate knotweed
Common buckwheat	Prostrate pigweed
Common chickweed	Redmaids
Common cocklebur*	Redroot pigweed
Common groundsel	Russian thistle*
Common lambquarters	Scentless chamomile/ mayweed
Common ragweed*	Shepherd's-purse
Common sunflower	Slimleaf lambquarters
Corn chamomile	Smallflower buttercup
Corn gromwell*	Smallseed falseflax
Corn spurry	Stinking chickweed
Cowcockle	Stinking mayweed/ dogfennel
Cress (<i>mouse-ear</i>)	Sunflower
Curly dock	Swinecress
False chamomile	Tansy mustard
Field chickweed	Tarweed fiddleneck
Field pennycress	Tumble/Jim hill mustard
Filaree (<i>redstem</i> , <i>Texas</i>)	Volunteer canola
Flixweed	Volunteer lentils
Green smartweed	Volunteer peas
Henbit	Wild buckwheat*
Kochia*	Wild chamomile
Ladysthumb	Wild mustard
Lanceleaf sage*	
London rocket	
Mallow (<i>common</i> , <i>little</i>)**	

Rapport™ BroadSpec

Herbicide

APPLICATION RECOMMENDATIONS

RATES

Wheat and barley: Apply 0.4 to 1 ounce per acre after the crop is in the 2-leaf stage, but before the flag leaf is visible.

Do not exceed 1 ounce per acre per crop season.

Fallow: Apply 0.4 to 1 ounce per acre to fallow in the spring, summer or fall when the majority of weeds have emerged and are actively growing.

Do not exceed 1 ounce per acre per crop season. Rapport BroadSpec should be applied in combination with other suitable registered fallow herbicides such as glyphosate plus 2,4-D (ester formulations work best) or glyphosate plus dicamba. When applied at a rate of 0.4 to 0.6 ounce per acre, Rapport BroadSpec must be used in a tank-mix combination with other registered fallow herbicides.

Pre-plant burndown: Apply 0.4 to 1 ounce per acre as a burndown treatment prior to, or shortly after planting (prior to emergence). The total amount cannot exceed 1 ounce per acre per crop season.

Apply up to 0.6 ounce per acre of Rapport BroadSpec as a pre-plant burndown to cotton. Allow at least 14 days from time of application to time of planting cotton.

Post harvest: Apply 0.4 to 1.0 ounce per acre to crop stubble after harvest. Use the 1.0 ounce per acre rate when weed infestation is heavy and consists mainly of those weeds listed under the "WEEDS PARTIALLY CONTROLLED" section of the label or when application timing and environmental conditions are marginal. Apply Rapport BroadSpec when all or most of the weeds have germinated.

Annual broadleaf weeds should be past the cotyledon stage, actively growing and less than 4" tall or wide. Rainfall immediately after treatment can wash Rapport BroadSpec off of weed foliage, resulting in reduced weed control. Several hours of dry weather are needed to allow Rapport BroadSpec to be sufficiently absorbed by weed foliage.

Better Choices. Better Business. | www.nufarm.com/us

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/us to download a full label.

©2010 Nufarm. Important: Always read and follow label instructions. Rapport™ is a trademark of Nufarm. Affinity® is a trademark of E.I. du Pont de Nemours and Company.

10-AG-0020

* See SPECIFIC WEED PROBLEMS for more information.
**Partial control

