



Rapport™ TankMix

Selective Herbicide

Rapport™ TankMix Selective Herbicide contains thifensulfuron and tribenuron, the active ingredients used in Affinity™ TankMix.



Rapport™ TankMix Makes Your Cereal Herbicide Program Better

Rapport™ TankMix Selective Herbicide delivers reliable control of a broad spectrum of broadleaf weeds for clean fields and maximum yield potential. The active ingredients in Rapport TankMix have been shown to be safe to growing crops while delivering dependable control of broadleaf weeds. Rapport TankMix boosts any weed-control program: When added to your small grains grass or kochia control program, Rapport TankMix will manage tough weeds like Canada thistle, lambsquarters, Russian thistle, dogfennel and wild buckwheat. Plus, with Rapport TankMix, you can rotate to almost any crop after harvest.

KEY PERFORMANCE BENEFITS

Dependable post-emergent control of a broad spectrum of broadleaf weeds
Low tank-mix antagonism
Excellent crop safety
Flexible rotation options

ACTIVE INGREDIENTS

thifensulfuron-methyl
and tribenuron-methyl

% ACTIVE INGREDIENTS

40% and 10%

FORMULATION

dry flowable

CHEMICAL FAMILY

sulfonylurea

EPA REGISTRATION

71368-80

BRAND ALTERNATIVE

Affinity™ TankMix

MODE OF ACTION

ALS inhibitor

GROUP NUMBERS

2

RESTRICTED USE

no

SIGNAL WORD

danger

PACKAGING INFORMATION

8 x 48 oz



WEEDS CONTROLLED

WHEN TANK MIXED WITH BROMOXYNIL	Smallflower buttercup	Miners lettuce	False chamomile	Common groundsel
Annual knawel	Smooth pigweed	Mouse-ear chickweed	Field pennycress	Common lambsquarters
Annual sowthistle	Spiny pigweed	Pennsylvania smartweed	Flixweed	Common mallow ^{†††}
Black mustard	Stinking mayweed/dogfennel	Pepperweed species	Fumitory	Common purslane ^{***}
Black nightshade	Swinecress	Prickly lettuce*‡	Giant ragweed	Common ragweed ^{***}
Bushy wallflower/treacle mustard	Tall morningglory	Prostrate knotweed	Green smartweed	Common sunflower ^{***}
Carolina geranium	Tall waterhemp	Puncturevine	Henbit	Corn chamomile
Coast fiddleneck	Tansy mustard	Redmaids	Ivyleaf morningglory	Corn spurry
Common buckwheat	Tartary buckwheat	Redroot pigweed	Kochia*‡	Cress (<i>mouse-ear</i>)
Common chickweed*	Tarweed fiddleneck	Russian thistle*‡	Ladysthumb	Curly dock
Common cocklebur	Tumble/Jim hill mustard	Scentless chamomile/mayweed	Marshelder	Cutleaf nightshade ^{†††}
Common groundsel	Velvetleaf	Shepherd's-purse	Miners lettuce	Eastern black nightshade ^{†††}
Common lambsquarters	Volunteer canola	Smallflower buttercup	Mouse-ear chickweed	False chamomile
Common mallow ^{**}	Volunteer lentils	Smooth pigweed	Pennsylvania smartweed	Field bindweed ^{†††}
Common ragweed	Volunteer peas	Spiny pigweed	Pepperweed species	Field horsetail ^{†††}
Common sunflower*	Wild buckwheat	Stinking mayweed/dogfennel	Prickly lettuce*‡	Field pennycress
Common tarweed	Wild chamomile	Swinecress	Prostrate knotweed	Green smartweed
Corn chamomile	Wild mustard	Tall morningglory [†]	Puncturevine	Hemp dogbane ^{***}
Corn gromwell	Wild radish	Tall waterhemp [†]	Redmaids	Henbit ^{†††}
Corn spurry	Yellow rocket	Tansy mustard	Redroot pigweed	Kochia*‡
Cowcockle	WHEN TANK MIXED WITH 2,4-D PRODUCTS (Such as Weedar® 64)	Tarweed fiddleneck	Russian thistle*‡	Ladysthumb
Cress (<i>mouse-ear</i>)	Annual knawel	Tumble/Jim hill mustard	Scentless chamomile/mayweed	London rocket
Cutleaf evening primrose ^{**}	Annual sowthistle	Velvetleaf	Shepherd's-purse	Marestail ^{†††}
Cutleaf nightshade	Black mustard	Volunteer canola	Smallflower buttercup	Mallow (<i>little</i>)
Curly dock	Bushy wallflower/treacle mustard	Volunteer lentils	Smooth pigweed	Marshelder
Eastern black nightshade	Carolina geranium	Volunteer peas	Spiny pigweed	Miners lettuce
False chamomile	Coast fiddleneck	Volunteer sunflower*	Stinking mayweed/dogfennel	Morningglory species ^{***}
Field pennycress	Common buckwheat	Wild buckwheat	Swinecress	Mouse-ear chickweed
Flixweed	Common cocklebur	Wild chamomile	Tall morningglory	Pennsylvania smartweed
Fumitory	Common groundsel	Coast fiddleneck	Tall waterhemp	Prickly lettuce ^{***‡}
Giant ragweed	Common lambsquarters	Common buckwheat	Tansy mustard	Prostrate knotweed
Green smartweed	Common mallow	Common cocklebur	Tarweed fiddleneck	Puncturevine ^{***}
Hemp sesbania	Common purselane	Common sunflower*	Tumble/Jim hill mustard	Redmaids
Henbit	Common ragweed	Common tarweed	Velvetleaf	Redroot pigweed
Horned poppy	Curly dock	Common sunflower*	Volunteer canola	Russian thistle*‡
Ivyleaf morningglory	False chamomile	Common tarweed	Volunteer lentils	Scentless chamomile/mayweed
Jimsonweed	Field pennycress	Common tarweed	Volunteer peas	Shepherd's-purse
Kochia*‡	Flixweed	Common tarweed	Volunteer sunflower*	Silverleaf nightshade ^{**}
Ladysthumb	Fumitory [†]	Common tarweed	Wild buckwheat	Silverleaf nightshade ^{**}
Lanceleaf sage	Giant ragweed	Common tarweed	Wild chamomile	Smallflower buttercup
London rocket	Green smartweed	Common tarweed	Wild mustard	Stinking mayweed/dogfennel
Mallow (<i>little</i>)	Hemp sesbania [†]	Common tarweed	Wild radish	Swinecress
Marestail ^{**}	Henbit	Common tarweed	WHEN TANK MIXED WITH FLUROXYPYR PRODUCTS	Tansy mustard
Marshelder	Ivyleaf morningglory	Common tarweed	Annual knawel	Tarweed fiddleneck
Miners lettuce	Kochia*‡	Common tarweed	Annual sowthistle	Tumble/Jim hill mustard
Mouse-ear chickweed	Ladysthumb	Common tarweed	Bedstraw (<i>cleavers</i>) ^{***}	Velvetleaf ^{***}
Pennsylvania smartweed	London rocket	Common tarweed	Black mustard	Venice mallow ^{***}
Pepperweed species	Mallow (<i>little</i>)	Common tarweed	Black nightshade ^{†††}	Volunteer canola
Prickly lettuce*‡	Marestail ^{††}	Common tarweed	Bushy wallflower/treacle mustard	Volunteer flax ^{***}
Prostrate knotweed	Marshelder	Common tarweed	Carolina geranium	Volunteer lentils
Puncturevine		Common tarweed	Coast fiddleneck	Volunteer peas
Redmaids		Common tarweed	Coffeeweed ^{***}	Volunteer potato ^{†††§}
Redroot pigweed		Common tarweed	Common buckwheat	Volunteer sunflower*
Russian thistle*‡		Common tarweed	Common chickweed ^{***}	Wild buckwheat
Scentless chamomile/mayweed		Common tarweed	Common cocklebur ^{***}	Wild chamomile
Shepherd's-purse		Common tarweed		Wild mustard
Silverleaf nightshade		Common tarweed		White clover ^{***}

* See "SPECIFIC WEED PROBLEMS" section of the Rapport TankMix label for more information.
^{**} Partial control: A visual reduction of weed population as well as a significant loss of vigor for individual weed plants. For better results, use 6 ounces active ingredient per acre of bromoxynil-containing herbicide.
[†] Partial control: A visual reduction of weed population as well as a significant loss of vigor for individual weed plants. For better results, use higher rates 2,4-D containing herbicides.
^{††} Partial control: A visual reduction of weed population as well as a significant loss of vigor for individual weed plants. For better results, use higher rates 2,4-D and/or dicamba-containing herbicides.
^{†††} Partial control: A visual reduction of weed population as well as a significant loss of vigor for individual weed plants. Use 1-1/2 - 2 ounces active ingredient per acre of fluroxypyr-containing herbicide.
[‡] Naturally-occurring resistant biotypes of kochia, prickly lettuce, and Russian thistle are known to occur. See the "TANK MIXTURES" and "SPECIFIC WEED PROBLEMS" sections of the label for additional details.
^{***} Use 1-1/2 - 2 ounces active ingredient per acre fluroxypyr containing herbicides.
[§] Use 2 - 4 ounces active ingredient per acre fluroxypyr-containing herbicides. See specific fluroxypyr-containing herbicide label for application rates and precautions.

Rapport™ TankMix

Selective Herbicide

LABELED USES

Wheat
 Barley
 Triticale
 Fallow
 Pre-plant or post-harvest burndown herbicide

APPLICATION RECOMMENDATIONS

APPLICATION INFORMATION

Rapport TankMix can be mixed with grass-control herbicides and other broadleaf herbicides for weed control in wheat, barley and/or triticale.

RATES

Apply 0.6 to 1.0 oz/A to select weeds from germination to 8 inches in height.

Wheat, barley and triticale: Make applications after the crop is in the 2-leaf stage, but before the flag leaf is visible.

Post-harvest: Apply when the majority of weeds have emerged and are actively growing.

Fallow: Apply when the majority of weeds have emerged and are actively growing.

Pre-Plant burndown: Apply up through planting, but before wheat, barley or triticale plants emerge. Apply at least 60 days before planting to sugarbeets, winter rape and canola fields. Apply at least 45 days before planting for any other crops (such as soybeans and field corn, cotton, rice or grain sorghum.)

Better Choices. Better Business.
www.nufarm.com/us

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/us to download a full label.

©2010 Nufarm. Important: Always read and follow label instructions. Rapport™ and Weedar® are trademarks of Nufarm. Affinity® is a trademark of E.I. du Pont de Nemours and Company.

10-AG-0021

