

**OUTLOOK CONSISTENTLY DELIVERS SEASON-LONG GRASS AND SMALL-SEEDED BROADLEAF WEED CONTROL IN A VARIETY OF WEATHER AND TILLAGE CONDITIONS.**

**ANNUAL GRASSES**

Barnyardgrass	Goosegrass	Sandbur <sup>1</sup>
Bluegrass, annual, roughstalk	Johnsongrass (seedling) <sup>2</sup>	Shattercane <sup>1</sup>
Brome, California, downy	Millet, wild proso <sup>1</sup>	Signalgrass, broadleaf <sup>1</sup>
Crabgrass, large, smooth	Panicum, fall, Texas <sup>1</sup>	Witchgrass
Cupgrass, southwestern, woolly <sup>1</sup>	Rice, red	
Foxtail, giant, green, yellow	Ryegrass, Italian	

**SEDGES**

Flatsedge, rice	Nutsedge, yellow <sup>2</sup>
-----------------	-------------------------------

**ANNUAL BROADLEAVES**

Amaranth, Palmer, Powell	Lambsquarters, common <sup>1</sup>	Purslane, common
Beggarweed, Florida <sup>1</sup>	Nightshade, <sup>2</sup> black, eastern	Pusley, Florida
Carpetweed	black, hairy, cutleaf	Ragweed, common <sup>1</sup>
Chamomile, mayweed	Pigweed, prostrate, redroot,	Spurge, nodding, spotted
Eclipta <sup>1</sup>	smooth, tumble	Waterhemp, common <sup>2</sup> , tall <sup>2</sup>

<sup>1</sup> Partial control or suppression. To complement control, **Outlook**<sup>®</sup> herbicide should be used in tank mixes or sequential applications with other herbicides that provide additional control of these weed species.

<sup>2</sup> For best control of these species, use the highest rate recommended by soil type. If dry conditions exist near application, or excessive rainfall occurs early in season, a postemergence herbicide or cultivation may be required to help control these weeds.

**For best results, use Outlook in a BASF-recommended, two-pass program followed by Distinct<sup>®</sup>, Marksman<sup>®</sup> or Clarity<sup>®</sup> herbicide. For more information, please visit [agproducts.basf.com](http://agproducts.basf.com).**



Always read and follow label directions.

Clarity, Distinct, Marksman and Outlook are registered trademarks of BASF. Dual II Magnum is a registered trademark of Syngenta Group Company.

Harness is a registered trademark of Monsanto Technology LLC. Surpass is a trademark of Dow Agrosciences LLC.

©2005 BASF Corporation. All Rights Reserved. APN 05-01-002-00010

# Outlook Preemergence Herbicide for Corn



## THE STRAIGHT GOODS HERBICIDE TO COUNT ON.

- **CONSISTENT GRASS AND BROADLEAF WEED CONTROL**
- **RELIABLE: SEASON-LONG RESIDUAL CONTROL FOR ALL TYPES OF WEATHER, TILLAGE AND ENVIRONMENTAL CONDITIONS**
- **LOWEST USE-RATES: 33% LESS THAN DUAL II MAGNUM<sup>®</sup>/50% LESS THAN HARNESS<sup>®</sup> AND SURPASS<sup>®</sup>**
- **WIDE APPLICATION WINDOW: CAN BE APPLIED IN FALL, AS WELL AS EARLY PRE-PLANT THROUGH EARLY POSTEMERGENCE**
- **EXCELLENT CROP SAFETY—NO SAFENERS FOR CORN REQUIRED/NO CROP ROTATION RESTRICTIONS**

### ■ SUPERIOR ACTIVITY. SUPERIOR CONTROL.

With its unique water-soluble chemistry, **Outlook**<sup>®</sup> herbicide provides greater activity in the weed germination zone for improved uptake and superior control of the toughest weeds like waterhemp, pigweed, nightshade and lambsquarters. From the start, **Outlook** requires just 1/4" of rain for activation. Yet, even under the wettest conditions, it binds more tightly to the soil to minimize leaching. For all tillage systems, including no-till, **Outlook** washes off residue more-readily. So more herbicide is available to control the toughest annual grasses and broadleaf weeds—all season long.

#### OUTLOOK



#### DUAL II MAGNUM



#### ACETOCHLOR (ACTIVE INGREDIENT IN HARNESS & SURPASS)



■ **SOIL DISPERSION STRING TEST SHOWS THE SUPERIOR ACTIVITY OF OUTLOOK IN THE WEED GERMINATION ZONE PROVIDING GREATER UPTAKE FOR MORE CONSISTENT PERFORMANCE.**

### LABELED CROPS

Corn (field, seed, popcorn, sweet), dry beans, peanuts, sorghum (grain), soybeans, and perennial grasses grown for seed, sugar beets, garlic, dry bulb onions, dry bulb shallots, horseradish, potato

### ACTIVE INGREDIENTS (MODE OF ACTION)

Dimethenamid-P (root and shoot growth inhibitor)

### APPLICATION METHODS AND TIMING

#### Preplant Surface (minimum or no-tillage)

Up to 45 days before planting

Notes: Use the highest rate recommended for the specific soil type. Not recommended for use on coarse-textured soils or in areas where average rainfall typically exceeds 40".

#### Preplant Incorporated

Up to 2 weeks before planting

Notes: Incorporate in upper 1" to 2" soil surface. Avoid deeper incorporation or reduced weed control may result.

#### Preemergence Surface

After planting, before crop emergence

Notes: Broadcast uniformly to the soil surface. Rainfall, sprinkler irrigation, or shallow mechanical incorporation after application is required to move into the upper soil surface where seeds germinate.

#### Early Postemergence

Up to 10-leaf (12") corn

Notes: Apply before weeds germinate or in a tank mix with products that control germinated weeds.

#### Lay-By Applications

Corn taller than 12" but less than 36"

Notes: Do not make lay-by applications of **Outlook**<sup>®</sup> herbicide to sweet corn.

### USE RATE PER ACRE

Soil Texture Coarse/Organic Matter Content	<3%	12-14 fluid oz./A	≥3% <sup>1</sup>	14-18 fluid oz./A
	Medium or Fine	14-18 fluid oz./A	18-21 fluid oz./A	

<sup>1</sup> The rates listed are intended for full-season control of targeted weeds.

<sup>2</sup> For all early preplant applications, use 21 fluid ounces of **Outlook** per acre.

