



*Trademark of Dow AgroSciences LLC

Fact Sheet

Preplant & preemergence broadleaf control in corn and soybeans

Product Advantages:

- Unique selectivity fits both corn and soybeans
- Foundation control of more than 30 broadleaf weeds
- Absorption occurs by both root and shoot uptake
- Packaged in pre-measured water soluble packets
- Can be tank-mixed with pre-plant or pre-emergent grass herbicides
- Works in minimum and conventional tillage
- Minimal rotation restrictions

Weed spectrum:

| | | | |
|------------------------|------------------------------|-----------------|--|
| Cocklebur ¹ | Mustard | Russian thistle | Venice Mallow |
| Kochia | Nightshades | Smartweed | Waterhemp |
| Lambsquarters | Ragweed, common ¹ | Sunflower | Wormwood, biennial ² |
| Pigweeds | Ragweed, giant ² | Velvetleaf | ¹ light-moderate infestations ² partial control |

Application timing:

- Pre-plant incorporated, pre-plant surface, pre-emergence; postemergence (corn-velvetleaf only)
- Apply up to 30 days before planting

Herbicide Components:

- 1 ounce (a.i.) flumetsulam at 1.25 ounce/acre rate
- (1.25 ounce/acre of Python is equal to flumetsulam provided in 2 pintst/acre of Broadstrike+Treflan)

Application instructions:

- 0.8 - 1.33 ounces per acre (consult label for recommended rates)
- Calculate number of packets needed for spray mix
- Packaging: 4 x 4 ounce pouches per bag (12.8 acres at 1.25 oz/A rate), 4 bags per box
- Check label for liquid fertilizer mixing or dry fertilizer impregnation instructions
- Apply in 10 to 40 gallons of carrier
- PPI applications-incorporate uniformly into top 2-3 inches of soil

Other Information:

- PPE: long pants, long sleeved shirt, shoes and socks, waterproof gloves
- Not a Restricted Use Product, REI is 12 hours
- Thoroughly clean spray equipment after use
- Check label for pH restrictions and soil insecticide interactions pertaining to corn
- Crop Rotational Interval:

| | |
|---|----------------------|
| corn, soybeans | 0 months |
| alfalfa, wheat, barley, dry beans, peas | 4 months |
| potatoes, grain sorghum | 12 months |
| sunflower | 18 months |
| sugar beets, canola | 26 months & bioassay |

This fact sheet is not a substitute for the product label.

