



# Corn Insect Control

**Stallion™, a foliar corn insecticide where unequalled broad spectrum control is needed.**



Cutworms



Stalkborer



Chinchbug



Japanese Beetle



Western Corn Rootworm



Northern Corn Rootworm



Western Bean Cutworm



Corn Earworm



Southwestern Corn Borer



European Corn Borer

## Application Recommendations

**3.75 - 11.75 oz/A** Cutworm, including Army Cutworm (other than Western Bean Cutworm)

**5 - 11.75 oz/A** Green Cloverworm; Meadow Spittlebug; Western Bean Cutworm

**9.25 - 11.75 oz/A** Aphids; Armyworm, fall; Armyworm, southern; Armyworm, true; Armyworm, yellowstriped ; Bean Leaf Beetle; Bird Cherry Oat Aphid; Cereal Leaf Beetle; Chinchbug; Common Stalkborer; Corn Borer, European; Corn Borer, southwestern; Corn Earworm; Corn Leaf Aphid; Corn Leafhopper; False Chinchbug; Flea Beetle; Grasshopper; Hopvine Borer; Hornworm; Japanese Beetle (adult); Sap Beetle (adult); Southern Leaf Beetle; Stalkborer; Stinkbug; Tobacco Budworm; Webworm

## **Special Information:**

- Use a minimum of 2 gallons of finished spray per acre by air or 15 gallons of finished spray per acre by ground.
- A maximum of 11.75 ounces of product may be applied per acre (0.025 lbs/acre zeta-cypermethrin + 0.25 lbs/acre chlorpyrifos) per application.
- A maximum of 35.25 ounces of product per acre per season including at-plant plus foliar applications (0.075 lbs/acre zeta-cypermethrin + 0.75 lbs/acre chlorpyrifos).
- Do not make more than three applications of any product containing chlorpyrifos per season, including the maximum allowed of two granular applications, at the 1 pound chlorpyrifos rate.
- If more than 1 pound active ingredient granular chlorpyrifos per acre is applied at-plant (for a maximum of 1.3 pound active ingredient per acre per season), only one additional application of a liquid product containing chlorpyrifos at 1 pound active ingredient per acre is allowed per season, for a total of 2.3 pounds chlorpyrifos per acre per season.
- Do not apply in tank mixes with Steadfast® or Lightning® herbicides.

**Stallion can be tank mixed with foliar fungicides and fertilizers.**

Always read and follow label directions. Stallion is a restricted use pesticide. FMC and Stallion are trademarks of FMC Corporation. Steadfast is a trademark of E.I. du Pont de Nemours & Co. Lightning is a trademark of BASF Agrochemical Products B.V. © 2011 FMC Corporation.

LC-08-03ER-01/11

1-888-59-FMC-AG • FMCcrop.com

**FMC**