

Soybeans & Weed Control

Fast Weed Control

Cobra[®] Herbicide controls over 54 problem weeds faster than other herbicides with no concerns to rotated crops. Additionally, tests have shown *Cobra* suppresses white mold.

- Fast effective weed control
- No carryover concerns
- White mold suppression (see other side)

Weed Efficiency Ratings (Postemergence)

			, ,		
Trade Name	Cobra 10 – 12.5 fl oz/A	Classic[®] 1/2 – 2/3 oz/A	FirstRate [®] 0.3 oz/A	Flexstar [®] 1 – 1.5 pts/A	Ultra Blazer [®] 1 – 1.5 pts/A
Active Ingredient	lactofen	chlorimuron	cloransulam	fomesafen	acifluorfen
Cocklebur	G	E	E	G	G
Jimsonweed	E	G	E	E	E
Morningglory, annual	F	F	G	G	G
Nightshades, annual	G	Р	Р	G	G
Pigweeds	E	G	Р	E	E
Ragweed, common	E	G	E	G	G
Ragweed, giant	G	F	E	G	F
Sunflower, common	F	E	E	F	F
Velvetleaf	F	G	G	G	Р
Waterhemps	Е	Р	Р	E	E
Excellent Good	d Fair	Poor			

How To Use – Weed Control

Rate	10 – 12.5 fl oz/A
Timing	Preemergence – R6 (full seed)
Spray volume	20 gal/A
Adjuvant	COC @ 1 – 2 pts/A or NIS @ 0.25% v/v based on temperature and relative humidity. Best results are obtained with COC @ 1 pt/A + NIS @ 0.25% v/v + 2 - 4 lbs AMS/A. See label for details. Valent recommends use of a Chemical Producers & Distributors Association-certified adjuvant.
PHI	45 days
REI	12 hours
Roundup Ready PLUS [®]	Cobra qualifies for Roundup Ready PLUS soybean incentives. See www.RoundupReadyPLUS.com for full details.




The Power of Cobra."

Soybeans & White Mold Suppression

Suppress White Mold

White mold is a common disease found in Midwestern soybean production that, depending on time of infection, can cause significant yield loss. University research has shown a postemergence tank mix of *Cobra* Herbicide applied to soybeans will suppress the development of white mold. Research indicates that after an application of *Cobra*, a soybean plant produces several isoflavones that serve to protect against disease development.

The Picture Says it All



White mold infestation 1% Cobra-treated 20% Untreated

Treated: Jul. 2 Photo Taken: Sep. 15

Poplar Grove, IL Field had a history of white mold

How To Use – White Mold Suppression

Rate	6 fl oz/A	
Timing	Postemergence, V5 – R2, but prior to infection	
Spray volume	20 gal/A	
Adjuvant	COC @ 1 – 2 pts/A or NIS @ 0.25% v/v based on temperature and relative humidity. See label for details. Valent recommends use of a Chemical Producers & Distributors Association-certified adjuvant.	
PHI	45 days	
REI	12 hours	



Products That Work, From People Who Care[®] | www.valent.com | 800-6-VALENT (682-5368) Always read and follow label instructions.

Cobra and Products That Work, From People Who Care are registered trademarks and The Power of Cobra is a trademark of Valent U.S.A. Corporation. Classic is a registered trademark of E.I. du Pont de Nemours and Company. FirstRate is a registered trademark of Dow AgroSciences, LLC. Flexstar is a registered trademark of Syngenta. Ultra Blazer is a registered trademark of United Phosphorus, Inc. Roundup Ready PLUS is a trademark of Monsanto Company. ©2013 Valent U.S.A. Corporation. 2014-COB-8002 9/13 mf