



#### Weeds Controlled:

- barnyardgrass
- Canada thistle¹
- cocklebur
- crabgrass
- fall panicum
- field sandbur
- foxtail species
- jimsonweed
- Johnsongrass, seedling
- kochia
- lambsquarters
- marestail
- morningglory, annual
- nightshade species
- pigweed species
- prickly sida
- · ragweed, common
- ragweed, giant
- shattercane
- smartweed
- sunflower
- velvetleaf
- Venice mallow
- waterhemp
- witchgrass
- woolly cupgrass
- yellow nutsedge

Burndown activity in minimum and no-till corn only

Partial list of weeds controlled, always refer to label

# **Designed for Use in Roundup Ready® Corn**

### 1st Pass - Soil Applied

SureStart® Herbicide Use Rates

Soil Texture	Soil OM < 3%	Soil OM ≥ 3%
Coarse	1.5 - 2.0 pt/A	1.5 - 2.0 pt/A
Medium	1.5 - 2.5 pt/A	1.75 - 3.0 pt/A
Fine	2.0 - 3.0 pt/A	2.0 - 3.0 pt/A

- Apply early preplant up to 30 days before planting, preplant incorporated, preemergence or to spike stage corn. For maximum residual, apply as close to planting as possible
- Effective in all tillage systems
- Three non-glyphosate modes of action for improved resistance management
- Provides 5-6 weeks of residual control, designed to be followed by a post application of glyphosate when weeds reach 4-6" in height
- Do not use with Counter insecticide; plant corn at least 1.5" deep
- Doesn't contain atrazine may be used in atrazine restricted areas

# 2nd Pass - Post Applied

- Apply Durango® DMA® herbicide at 24 oz/A to 4-6" weeds
- Optimal weed control timing for maximum yield protection

# **Early Post Tankmix**

- SureStart® herbicide + Durango DMA tank mixture may be applied from spike to 11" tall corn
- Apply to weeds 1–2" in height
- Do not use liquid fertilizer as the carrier

SureStart® 1.75 – 2.0 pt/acre
Durango® DMA® 24 oz/acre
AMS 2.5 lbs/acre

**NOTES:** 

Science, Yield, Success.™

