

Trooper™ P+D

SELECTIVE HERBICIDE

Two-in-One Formula for Fast and Lasting Control.

Trooper™ P+D is a convenient premix of picloram and 2,4-D that works fast with quick burndown and provides season-long control through soil residual. It controls many annual and perennial weeds, prevents encroaching brush, and is safe to desirable grasses. Trooper P+D is labeled for use on rangeland and permanent grass pasture, fallow cropland, non-cropland and CRP – without grazing restrictions – making it an excellent weed control choice.

COMPLETE CONTROL

- Convenient premix of picloram and 2,4-D
- Translocates throughout the plant for complete control to the roots
- Provides lasting weed control through soil residual

EXCELLENT FOR RANGE AND PASTURE

- No grazing restrictions, except for lactating dairy animals (see label for specific restrictions)
- Safe to most cool- and warm-season grasses

EPA REG. NO.	228-530
ACTIVE INGR.	picloram <i>(10.2%)</i> , 2,4-D <i>(39.6)</i>
FORMULATION	liquid
CHEM. FAMILY	pyridines, phenoxy acetic acids
GROUP NO.	4
MODE OF ACT.	•
	synthetic auxin
MODE OF ACT.	synthetic auxin
MODE OF ACT. SIGNAL WORD RESTRICTED	synthetic auxin warning



Trooper™ P+D

SELECTIVE HERBICIDE

KEY USES

Rangeland & permanent grass pastures Conservation Reserve Program (CRP)

KEY WEEDS CONTROLLED

(See product label for complete list)

Alder

Aspen

Bindweed, field

Birch

Blackberry

Bouncingbet

Buttonbrush

Carrot, wild

Cherry

Chicory

Clover

Dandelion

Dock

Elm

Fleabane

Goldenrod

Hemlock

Hickory

Honeysuckle

Horsenettle

Knapweed

Kudzu

Locust

Maple Oak

Persimmon

Plantain

Poison oak

Prickly lettuce

Ragweed

Ragwort, tansy

Rush skeleton weed

Sassafras

Sourwood

Sowthistle

Spurge, leafy

Starthistle, yellow

Thistles

Toadflax

Tulip poplar

Vetch

Willow

APPLICATION RECOMMENDATIONS

BROADCAST FOLIAR APPLICATION (Ground or Aerial): Unless otherwise specified, apply in water alone or in an oil-water emulsion in a total spray volume of 10 to 40 gallons per acre using ground equipment or 1 or more gallons per acre by aerial application. If aerially applied, results will be more consistent for spray volumes of 2 or more gallons per acre. Use of the lower total spray volume with ground equipment is recommended primarily where this product is applied simultaneously with liquid fertilizer. Good coverage is essential. For aerial application, swath width should not exceed 1-1/4 times the wingspan of the aircraft.

To provide more complete wetting and coverage of the foliage, a non-ionic surfactant may be used at recommended rates. The use of a drift control additive is recommended for drift reduction and improved deposition.

See label for complete application recommendations.

RATES: Broadleaf weeds and woody plants in rangeland and permanent grass pastures SOUTHWEST, SOUTHEAST, AND MID-ATLANTIC STATES:

2-4 pints/acre depending on weed specie

NORTH AND NORTHWESTERN U.S.: (Colorado, Idaho, Iowa, Kansas, Minnesota, Missouri, Montana, Nebraska, North Dakota, Oregon, South Dakota, Utah, Washington and Wyoming) 2-7.5 pints/acre depending on weed specie

See label for complete application rates.

PASTURE AND RANGELAND USE RESTRICTIONS

- Maximum single application rate is 7.5 pints of product (0.5 lbs picloram ae) per acre for both broadleaf and woody weed control
- Maximum yearly application rate is 15 pints of product (1.0 lbs picloram ae) per acre
- Spot treatments and broadcast treatments can be applied during the same growing season only if the total amount of product applied does not exceed 15 pints (1.0 lbs picloram ae) per acre
- Spot treatment areas are defined as areas less than 1,000 sq ft

GRAZING RESTRICTIONS

- There are no grazing restrictions for non-lactating dairy animals or other livestock including horses, sheep, goats, and other animals in the treatment area
- Do not allow lactating dairy animals to graze treated areas within 7 days after application
- Do not harvest grass cut for hay from treated areas for 30 days after application
- Meat animals must be withdrawn from treated forage at least 3 days before slaughter



