

**CLARITY HERBICIDE CONTROLS OR SUPPRESSES MORE
THAN 190 ANNUAL AND PERENNIAL BROADLEAF WEEDS.**

ANNUAL WEEDS CONTROLLED

Amaranth, Palmer	Ladysthumb	Sicklepod
Beggarweed, Florida	Lambsquarters, common	Sida, prickly (teaweed)
Buckwheat, wild	Mallow, Venice	Smartweed, Pennsylvania
Buffalobur	Marestail	Sowthistle, annual
Burcucumber	Morningglory	Spurge, prostrate
Carpetweed	Nightshade	Sunflower, wild, volunteer
Cocklebur, common	Pigweeds	Thistle, Russian
Croton, tropic	Purslane, common	Velvetleaf
Jimsonweed	Ragweed, common, giant	Waterhemp, common, tall
Knotweed, prostrate	Sesbania, hemp	
Kochia	Shepherdspurse	

PERENNIAL AND BIENNIAL WEEDS SUPPRESSED

Alfalfa	Horsenettle, Carolina	Thistle, Canada
Bindweed, field, hedge	Milkweed, common, honeyvine	Burdock
Dandelion	Nightshade, silverleaf	Sweetclover
Dock, broadleaf, curly	Smartweed, swamp	Thistle, bull, milk, musk, plumeless
Dogbane, hemp	Sowthistle, perennial	

**For best results, use Clarity® herbicide in a BASF-recommended,
sequential program following Outlook® herbicide.
For more information, please visit agproducts.basf.com.**

Clarity[®]
herbicide

Always read and follow label directions.

Clarity and Outlook are registered trademarks of BASF.
©2005 BASF Corporation. All Rights Reserved. APN 05-01-002-0001C

Clarity Postemergence Herbicide for Corn



BASF
The Chemical Company

POST PARTNER FOR MORE CONSISTENT BROADLEAF WEED CONTROL.

- CONTROLS OR SUPPRESSES MORE THAN 190 ANNUAL AND PERENNIAL BROADLEAF WEEDS
- STOPS EVEN THE TOUGHEST ALS- AND ATRAZINE-RESISTANT SPECIES LIKE WATERHEMP AND KOCHIA
- A SINGLE PASS PROVIDES QUICK KNOCKDOWN
- WIDE WINDOW OF APPLICATION – FROM 5 TO 7 DAYS AFTER PLANTING, UP TO 36-INCH-TALL CORN
- CONSISTENT: WORKS IN VARIABLE TILLAGE AND WEATHER CONDITIONS, REQUIRES JUST 1/4" OF WATER FOR ACTIVATION
- EXCELLENT CROP SAFETY
- LOW VAPOR DRIFT – 8 TIMES LESS VOLATILE THAN GENERIC DMA DICAMBA SALTS




VERSATILE, CONSISTENT, SEASON-LONG CONTROL WITHOUT CARRYOVER.

With **Clarity**® herbicide, you have the versatility to deal with unpredictable spring weather. Its application window extends from as early as 5 to 7 days after planting, until 36" tall corn – giving you weeks to spray. And it provides consistent, economic control of the toughest broadleaf weeds like waterhemp and kochia. With quick knockdown of problem weeds in all types of weather and tillage conditions, **Clarity** is effective on weeds and safe for corn.

Lower use rates than other dicamba products along with lower volatilization rates helps **Clarity** stay on the leaf surface.

WIDE WINDOW OF APPLICATION:



LABELED CROPS

Corn (field, pop and seed), grain sorghum and fallow systems

ACTIVE INGREDIENTS (MODE OF ACTION)

Diglycolamine (Dicamba) – a growth regulator, Dicamba interferes with the plant’s growth hormones (auxins).

APPLICATION METHODS AND TIMING

- Preplant burndown** Prior to planting
Notes: Apply **Clarity**® herbicide at least 15 days prior to sorghum planting.
- Preemergence** After planting but prior to corn emergence
Notes: When planting into a legume sod (e.g., alfalfa or clover) in no tillage corn, apply **Clarity** after 4" to 6" of regrowth has occurred. In conventional or reduced tillage corn, do not apply to coarse-textured soils (sand, loamy sand, or sandy loam) or any soil with less than 2.5% organic matter until after corn emergence. **Clarity** is not labeled for a preemergence application in sorghum.
- Early postemergence** After corn emergence, up to 5-leaf or 8" tall corn
Notes: In sorghum, apply **Clarity** in the spike stage (all sorghum emerged) but before sorghum is 15" tall. For best performance, apply **Clarity** when the sorghum crop is in the 3- to 5-leaf stage and weeds are small (less than 3" tall). Use drop nozzles if sorghum is taller than 8".
- Late postemergence** From 8" to 36" tall corn or 15 days before tassel emergence, whichever first
Notes: For best performance, apply when weeds are less than 3" tall. Apply directed spray when corn leaves prevent proper spray coverage, sensitive crops are growing nearby, or tank mixing with 2,4-D. Do not apply **Clarity** when soybeans are growing nearby if any of these conditions exist: corn is more than 24" tall; soybeans are more than 10" tall; soybeans have begun to bloom.

USE RATE PER ACRE

Soil Texture Coarse/Organic Matter Content	≥2.5% ¹	8 fl. oz./A
	Medium or Fine	16 fl. oz./A ²

¹ Use 8 fluid ounces of **Clarity** per acre on soils with less than 2.5% organic matter.

² Apply only 8 fluid ounces of **Clarity** per acre for late postemergence.

