

**THE KNOCKDOWN POWER OF GLYPHOSATE, PLUS
THE RESIDUAL CONTROL OF PURSUIT.**

ANNUAL GRASSES CONTROLLED

Barley, volunteer	Johnsongrass, seedling, rhizome	Signalgrass, broadleaf
Barnyardgrass	Oats, volunteer	Sorghum, alumum
Crabgrass, large, smooth	Panicum, fall, Texas, browntop	Sprangletop
Cupgrass, woolly	Rice, red	Wheat, volunteer
Foxtail, giant, green, yellow	Rye	Wild oats
Goosegrass	Shattercane	

SEDGES (REDUCED COMPETITION)

Nutsedge, purple, yellow

ANNUAL BROADLEAF WEEDS CONTROLLED

Alligatorweed	Fleabane, annual, hairy, rough	Pigweed, redroot, smooth, spiny
Amaranth, Palmer	Jimsonweed	Ragweed, common, giant
Anoda, spurred	Hemp sesbania	Sicklepod
Artichoke, Jerusalem	Horseweed/marestail	Smartweed, ladysthumb, Pennsylvania
Buffalobur	Knotweed	Spurge, prostrate, spotted
Buckwheat, wild	Kochia	Starbur, bristly
Burcucumber	Lambsquarters, common	Sunflower
Carpetweed	Marshelder	Teaweed/prickly sida
Chickweed	Morningglory, annual	Velvetleaf
Cocklebur, common	Mustards	Waterhemp
Copperleaf, hophornbeam, Virginia	Nightshade, black, eastern black, hairy	
Eclipta		

For best results in Roundup Ready soybeans, use **Extreme®** herbicide postemergence in a BASF-recommended, two-pass program following **Prowl® H₂O** herbicide at burndown. For more information, please visit agproducts.basf.com.



Always read and follow label directions.

Extreme, Prowl and Pursuit are registered trademarks of BASF.
Roundup and Roundup Ready are registered trademarks of Monsanto Technology LLC.
©2005 BASF Corporation. All Rights Reserved. APN 05-01-002-0001Ex

Extreme Herbicide for Roundup Ready Soybeans and No-till



BASF
The Chemical Company

BURNDOWN WEED CONTROL IN NO-TILL SOYBEANS AND STALE SEEDBED

- **2-IN-1 EQUALS MORE CONTROL** – **EXTREME®** HERBICIDE PROVIDES THE QUICK KNOCKDOWN POWER OF GLYPHOSATE PLUS THE PROVEN RESIDUAL CONTROL OF **PURSUIT®** HERBICIDE
- **EXTREMELY EFFECTIVE AGAINST DIFFICULT BROADLEAF AND GRASSY WEEDS FOUND IN NO-TILL, SUCH AS MARESTAIL**
- **TANK MIXES EASILY FOR BURNDOWN** – WITH **PROWL® H₂O** HERBICIDE FOR SEASON-LONG CONTROL OF GRASSES AND ENHANCED CONTROL OF WATERHEMP, OR WITH 2,4-D TO CONTROL HEMP DOGBANE
- **FLEXIBILITY: CAN BE APPLIED FOR BURNDOWN, AT PREPLANT, OR AT PREEMERGENCE IN ANY SOYBEAN VARIETY RESIDUAL CONTROL**

POSTEMERGENCE WEED CONTROL IN ROUNDUP READY WITH RESIDUAL

- **SAFE FOR POSTEMERGENCE APPLICATION ON ROUNDUP READY® (GLYPHOSATE-TOLERANT) SOYBEANS ONLY!** – WHEN WEEDS ARE ACTIVE AND GROWING UNTIL 1" to 8" TALL
- **CLEANS-UP TOUGHEST WEEDS THAT CAN ESCAPE PRE-PLANT APPLICATIONS OF HERBICIDE IN ROUNDUP READY SOYBEANS**
- **RESIDUAL CONTROL OF GRASSY WEEDS AND BROADLEAVES**
- **RAINFAST AFTER ONE HOUR, EFFECTIVE EVEN IN COOLER-THAN-NORMAL WEATHER**

PART GLYPHOSATE. PART PURSUIT. ONE PROMISE.

Extreme is the leader of a category of value-added herbicides created specifically for maximum weed control in Roundup Ready soybeans. **Extreme** attacks newly emerged grasses and broadleaf weeds with the knockdown power of glyphosate and keeps them down with the trusted residual control of **Pursuit**, while clearing Roundup Ready soybeans of early-season, yield-robbing weed competition.

LABELED CROPS

Roundup Ready soybeans (postemergence control)
No-till/stale seedbed soybeans (burndown weed control)

ACTIVE INGREDIENTS (MODE OF ACTION)

Imazethapyr (ALS/AHAS enzyme inhibitor that disrupts protein synthesis and inhibits growth)
Glyphosate (EPSP enzyme inhibitor that disrupts amino acid synthesis)

APPLICATION METHODS AND TIMING

No-till, stale seedbed soybeans	Apply as burndown prior to planting Apply as preemergence of any soybean variety
Roundup Ready soybeans	Apply in-crop as first glyphosate on the acre*

*Notes: On Roundup Ready soybeans only, apply when weeds are actively growing and between 1" to 8" tall. Should be applied a minimum of one hour before rainfall or overhead irrigation.

USE RATES PER ACRE

3 pints per acre*

*Notes: Only one (1) application of **Extreme®** herbicide may be made during the season. When tank-mixed, always use in accordance with the more restrictive label limitations and precautions.

Adjuvants: Burndown or postemergence applications of **Extreme** require the addition of a surfactant and fertilizer. See label for other specific details.

BETTER RESIDUAL CONTROL WITH EXTREME



EXTREME HERBICIDE
8 WEEKS AFTER BURNDOWN.



ROUNDUP® HERBICIDE
8 WEEKS AFTER BURNDOWN.

In side-by-side 2001 field trials at the BASF Research Farm in Seymour, Illinois, the results are clearly superior with **Extreme**.

