

FirstRate[®]

PRE- AND POSTEMERGENCE CONTROL OF KEY BROADLEAF WEEDS

HERBICIDE

Product Advantages

- Provides both pre- and postemergence control of many broadleaf weeds, including glyphosate-resistant species
- Easily tank-mixed with many pre- and post emergence grass and broadleaf herbicides
- Effective in all tillage systems
- Excellent crop safety

Crop Rotation Interval	
Wheat	4 months
Alfalfa, corn, cotton, sorghum, rice, dry beans, oats, peanuts	9 months
Barley	12 months
Potatoes, sweet corn	18 months
Sugar beets, sunflowers	30 months & bioassay

Refer to the FirstRate® herbicide label for additional crop rotation interval restrictions.

Application Timing



• Preplant: Apply up to two weeks prior to planting or preemergence for additional burndown and early season residual control.

Rate: 0.3 to 0.6 ounce per acre

• Postemergence: Tank-mix with glyphosate to attack and prevent hard-to-control and glyphosate-resistant weeds. FirstRate may be applied up to the 50% flowering stage of soybeans.

Rate: 0.3 to 0.6 ounce per acre (recommended rate of 0.45 ounce)

Control Target Weeds



Lambsquarters



Marestail





Pigweed



Morningglory

Common Ragweed

Weed Spectrum

Burcucumber ¹	
Cocklebur	
Lambsquarters ²	
Mallow, Venice	
Marestail	
Marshelder ³	

Morningglory² Mustard, wild³ Nutsedge, yellow¹ Pigweed² Ragweed, common Ragweed, giant Smartweed, Pennsylvania Sunflower Thistle, Canada¹ Velvetleaf

> ¹ Suppression ² Soil-applied ³ Post-applied

Load Chart for FirstRate® herbicide

	Rate per acre						
Acres	0.3	3 oz.	0.45 oz.		0.6 oz.		
per load	WSP	Bags	WSP	Bags	WSP	Bags	
40	20	2	30	3	40	4	
80	40	4	60	6	80	8	
160	80	8	120	12	160	16	

Note: Each water soluble packet (WSP) is 0.6 ounce. There are 10 WSP per bag. Each box of FirstRate® herbicide contains four bags or 40 WSP.

For more information on FirstRate® herbicide, visit www.dowagro.com/usag.